

these and other nucleic acids of the present invention may be used as molecular weight standards in, for example, gel electrophoresis.

The term "cDNA" is intended to refer to DNA prepared using messenger RNA (mRNA) as template. The advantage of using a cDNA, as opposed to genomic DNA or DNA polymerized from a genomic, non- or partially-processed RNA template, is that the cDNA primarily contains coding sequences of the corresponding protein. There may be times when the full or partial genomic sequence is preferred, such as where the non-coding regions are required for optimal expression or where non-coding regions such as introns are to be targeted in an antisense strategy.

It also is contemplated that TIE-2, EDNRA, TGF β 3, TGF β III, VEGFR1, VEGF or bFGFR may be represented by natural variants that have slightly different nucleic acid sequences but, nonetheless, encode the same proteins (see Table 1 below).

As used in this application, the term "a nucleic acid encoding a TIE-2, EDNRA, TGF β 3, TGF β III, VEGFR1, VEGF or bFGFR" refers to a nucleic acid molecule that has been isolated free of total cellular nucleic acid. The term "functionally equivalent codon" is used herein to refer to codons that encode the same amino acid, such as the six codons for arginine or serine (Table 1, below), and also refers to codons that encode biologically equivalent amino acids, as discussed in the following pages.

Table 1

Amino Acids			Codons			
Alanine	Ala	A	GCA	GCC	GCG	GCU
Cysteine	Cys	C	UGC	UGU		
Aspartic acid	Asp	D	GAC	GAU		
Glutamic acid	Glu	E	GAA	GAG		
Phenylalanine	Phe	F	UUC	UUU		
Glycine	Gly	G	GGA	GGC	GGG	GGU
Histidine	His	H	CAC	CAU		

TABLE 1 - Continued

Isoleucine	Ile	I	AUA	AUC	AUU			
Lysine	Lys	K	AAA	AAG				
Leucine	Leu	L	UUA	UUG	CUA	CUC	CUG	
				CUU				
Methionine	Met	M	AUG					
Asparagine	Asn	N	AAC	AAU				
Proline	Pro	P	CCA	CCC	CCG	CCU		
Glutamine	Gln	Q	CAA	CAG				
Arginine	Arg	R	AGA	AGG	CGA	CGC	CGG	
				CGU				
Serine	Ser	S	AGC	AGU	UCA	UCC	UCG	
				UCU				
Threonine	Thr	T	ACA	ACC	ACG	ACU		
Valine	Val	V	GUA	GUC	GUG	GUU		
Tryptophan	Trp	W	UGG					
Tyrosine	Tyr	Y	UAC	UAU				

- The DNA segments of the present invention include those encoding biologically functional equivalent TIE-2, EDNRA, TGF β 3, TGF β RIII, VEGFR1, VEGF or bFGFR proteins and peptides, as described above. Such sequences may arise as a consequence of codon redundancy and amino acid functional equivalency that are known to occur naturally within nucleic acid sequences and the proteins thus encoded. Alternatively, functionally equivalent proteins or peptides may be created *via* the application of recombinant DNA technology, in which changes in the protein structure may be engineered, based on considerations of the properties of the amino acids being exchanged. Changes designed by man may be introduced through the application of site-directed mutagenesis techniques or may be introduced randomly and screened later for the desired function, as described below.